

Homework 1. Due Tuesday, January 27

Problem 1. *From Classical Mechanics by J. R. Tylor*

Problems: 5.13

Problem 2. *From Classical Mechanics by J. R. Tylor*

Problems: 7.30


Homework 2. Due Tuesday, February 3

Problem 1. 1999-Spring-CM-U-3

A classical oscillator consisting of two masses m_1 and m_2 , connected by an ideal spring (spring constant k), slides on a frictionless ramp as shown. The lower portion of the ramp is horizontal (gravity acts vertically), while on the left is an immovable vertical wall. The motion of the oscillator occurs entirely within the plane of the figure. The system starts stationary at height h , with the spring un-stretched.

Assume collisions with the wall, of which there may be more than one, are instantaneous and elastic. Neglect the small height difference between masses, and also neglect any differential force on them caused by the curvature of the ramp. Furthermore, assume the spring is sufficiently stiff that it does not collapse, so that the masses never touch each other.

1. For m_1 sufficiently greater than m_2 , there will be only one collision with the wall before the oscillator starts back up the ramp. In this case, find the maximum height attained by the oscillator on its first rebound.
2. For $m_1 = m_2 = m$, determine the behavior of the oscillator as it interacts with the wall. Find the maximum height attained by the rebounding oscillator in this situation, and sketch x_1 and x_2 vs. time while the oscillator interacts with the wall.
3. In the case that $m_1 \ll m_2$, describe qualitatively the behavior while the oscillator interacts with the wall, and give qualitative sketches of x_1 , and x_2 vs. time, demonstrating this behavior.



Problem 2. 2001-Spring-CM-U-3


A particle with mass m , confined to a plane, is subject to a force field given by

$$\vec{F}(\vec{r}) = \begin{cases} -k\vec{r}, & \text{for } |r| \leq a \\ 0, & \text{for } |r| > a \end{cases}$$

The particle enters the field from the left horizontally with an initial velocity v_0 and a distance $b < a$ below the force center as shown.

1. Write down the equation of motion for $r \leq a$ in Cartesian coordinates in terms of x , y , and $\omega = \sqrt{k/m}$

2. Give the trajectory of the particle when $r < a$.
3. For $v_0 = a\omega$ find the coordinates of the particle as it exits the region of non-zero force field.
4. For $v_0 = a\omega$, find the deflection angle θ of the departing velocity at the exit point.



Homework 3. Due Tuesday, February 10

Problem 1. 1989-Spring-CM-U-1.

A particle of mass m scatters off a second particle with mass M according to a potential

$$U(r) = \frac{\alpha}{r^2}, \quad \alpha > 0$$

Initially m has a velocity v_0 and approaches M with an impact parameter b . Assume $m \ll M$, so that M can be considered to remain at rest during the collision.

1. Find the distance of closest approach of m to M .
2. Find the laboratory scattering angle. (Remember that M remains at rest.)

Problem 2. 1998-Spring-CM-U-3


A particle of mass m has velocity v_0 when it is far from the center of a potential of the form:

$$V(r) = \frac{k}{r^4},$$

where k is a positive constant that you can conveniently write as

$$k = \frac{1}{2}mv_0^2a^4.$$

1. Find r_{min} the distance of the closest approach of the particle to the center of force, as a function of the impact parameter b of the particle.
2. Find an expression for the scattering angle of the particle as a function of the impact parameter b . Your expression may include a definite integral that you need not evaluate, so long as no unknown quantities appear in the integral.



For Honors.

Problem 3. 1994-Fall-CM-G-3.jpg

A particle of mass m moves under the influence of an attractive central force $F(r) = -k/r^3$, $k > 0$. Far from the center of force, the particle has a kinetic energy E .

1. Find the values of the impact parameter b for which the particle reaches $r = 0$.
2. Assume that the initial conditions are such that the particle misses $r = 0$. Solve for the scattering angle θ_s , as a function of E and the impact parameter b .

Homework 4. Due Tuesday, February 17

Problem 1.

Determine the cross-section for a particle of energy E to “fall” to the center of a field $U = -\frac{\alpha}{r^2}$, where $\alpha > 0$.


Problem 2.

Determine the cross-section for a particle of energy E to “fall” to the center of a field $U = -\frac{\alpha}{r^n}$, where $\alpha > 0$ and $n > 2$.

Homework 5. Due Tuesday, February 24

Problem 1. 1983-Fall-CM-U-2.

Take $K = 4k$ and $m_1 = m_2 = M$. At $t = 0$ both masses are at their equilibrium positions, m_1 has a velocity \vec{v}_0 to the right, and m_2 is at rest. Determine the distance, x_1 , of m_1 from its equilibrium position at time $t = \frac{\pi}{4} \sqrt{\frac{m}{k}}$. Hint: First find the normal modes and the normal mode frequencies, then put in the initial conditions.



Problem 2.

Determine the normal/eigen frequencies for the four equal masses m on a ring. The masses are connected by the identical springs of spring constant k along the ring.

For Honors.

Problem 3.

Determine the normal/eigen frequencies for two equal masses m on a ring of radius R . The masses are connected by the identical springs of spring constant k along the ring. The ring is in the vertical plane. The acceleration of the free fall is g . Consider only the case of hard springs: $\frac{mg}{kR} \ll 1$.

Homework 6. Due Tuesday, March 3

Problem 1. 1985-Fall-CM-U-2.


A straight rod of length b and weight W is composed of two pieces of equal length and cross section joined end-to-end. The densities of the two pieces are 9 and 1. The rod is placed in a smooth, fixed hemispherical bowl of radius R . ($b < 2R$).

1. Find expression for the fixed angle β between the rod and the radius shown in Fig.1
2. Find the position of the center of mass when the rod is horizontal with its denser side on the left (Fig. 1). Give your answer as a distance from the left end.
3. Show that the angle θ which the rod makes with the horizontal when it is in equilibrium (Fig. 2) satisfies

$$\tan \theta = \frac{2}{5} \frac{1}{\sqrt{(2R/b)^2 - 1}}$$

Note the fundamental principles you employ in this proof.



4. Show that the equilibrium is small under stable displacements.



Problem 2. 1990-Fall-CM-U-1.


A small steel ball with mass m is originally held in place by hand and is connected to two identical horizontal springs fixed to walls as shown in the left figure. The two springs are **unstretched** with natural length L and spring constant k . If the ball is now let go, it will begin to drop and when it is at a distance y below its original position each spring will stretch by an amount x as shown in the right figure. It is observed that the amount of stretching x is very small in comparison to L .

1. Write down the equation that determines $y(t)$. (Take y to be positive in going downward.) Is this simple harmonic motion?
2. Find the equilibrium position y_{eq} about which the steel ball will oscillate in terms of m , g , k , and L .
3. Find the maximum distance, y_{max} , that the steel ball can drop below its original position in terms of m , g , k , and L .
4. Write down an expression for the period of the steel ball's motion. (DO NOT evaluate the integral.)



Homework 7. Due Tuesday, March 10

Problem 1.




Find normal frequencies and normal modes of a double pendulum.

Problem 2.

Three infinite horizontal rails are at distance l from each other. Three beads of equal masses m can slide along the rails without friction. The three beads are connected by identical springs of constants k and equilibrium/unstretched length $l/2$.


1. Find the normal modes and the normal frequencies.
2. At time $t = 0$ one of the masses is suddenly given a velocity v . What will be the velocity of the whole system of three masses (average velocity) in the long time?



Homework 8. Due Tuesday, March 17


Problem 1. *Three pendulums*

Three identical pendulums are connected by the springs as shown on the figure. Find the normal modes



Problem 2. *A pendulum a spring and a block*


A ball of mass m is hanging on a weightless spring of spring constant k , which is attached to a block of mass M . The block can move along the table without friction. The length of the unstretched spring is l_0 . Find the normal modes of small oscillations.



Homework 9. Due Tuesday, March 24


Problem 1. *Cylinders and a pendulum*

A uniform solid cylinder of radius r and mass m can roll inside a hollow cylinder of radius $R > r$ without slipping. A pendulum of length $l = \frac{R-r}{2}$ and mass $M = m/2$ is attached to the center of the smaller cylinder. Find the normal frequencies and normal modes of this system.



Problem 2. *A rod and a pendulum*

A uniform rod AB of mass M and length $3a$ can rotate around a point O , $|AO| = a$. A pendulum of length $l = a$ and mass $m = \frac{3}{4}M$ is attached to the end A of the rod. Find the normal frequencies and normal modes.



Homework 10. Due Tuesday, March 31

Problem 1.

1. Find the principal moments of inertia for a uniform square of mass M and side a .
2. Find the kinetic energy of the uniform cube of mass M and side a rotating with angular velocity Ω around its large diagonal.


Problem 2.

1. Find the kinetic energy of the uniform circular cone of height h , base radius R , and mass M .
2. The same for a uniform ellipsoid of semiaxes a , b , and c .

Homework 11. Due Tuesday, April 7

Problem 1. *A coin*

A uniform thin disc of mass m rolls without slipping on a horizontal plane. The disc makes an angle α with the plane, the center of the disc C moves around the horizontal circle with a constant speed v . The axis of symmetry of the disc CO intersects the horizontal plane in the point O . The point O is stationary. Find the kinetic energy of the disc.




Problem 2.

Ignoring interaction between the Earth and both the Moon and the Sun, find the precession angular velocity of the Earth. Consider Earth to be a uniform spheroid with semi-axes S and A and $\frac{C-A}{A} \approx \frac{1}{300}$.

For Honors.

Problem 3.

A cylindrical cone of mass M , height h , and angle α is rolling on the plane without slipping. Find its kinetic energy as a function of θ .



Homework 12. Due Tuesday, April 14

Problem 1. *A parallelepiped*


A uniform parallelepiped of mass m and edges a , b , and c is rotating with the constant angular velocity ω around an axis which coincides with the parallelepiped's large diagonal.

1. What is the parallelepiped's kinetic energy?
2. What torque must be applied to the axes in order to keep it still? (neglect the gravity.)

Problem 2. *A disk in a frame*

A square weightless frame $ABCD$ with the side a is rotating around the side AB with the constant angular velocity ω_1 . A uniform disk of radius R and mass m is rotating around the frame's diagonal AC with the constant angular velocity ω_2 . The center of the disk coincides with the center of the frame. Assuming $\omega_1 = \omega_2 = \omega$ find


1. The kinetic energy of the system.
2. The magnitude of the torque that needs to be applied to the axes to keep it still. (neglect the gravity.)



Homework 13. Due Tuesday, April 21

Problem 1.

A parallelepiped with edges a , b , c was deformed as shown on the figure. Find the strain tensor in the linear approximation assuming that θ is small. Remember, the answer must be a 3×3 tensor.



Problem 2.

For the deformation from the previous problem assuming that the shear modulus is μ find:

1. The stress tensor.
2. The force F which has to be applied to the top surface.
3. Total torque of all external forces.
4. Are you sure your answer for the previous question is correct?
5. How come the parallelepiped is not rotating then?

Homework 14. Due Tuesday, April 28

Problem 1.

A cylinder of radius R and length L is squeezed by applying a uniform pressure P_0 to its end.

1. Find the change of length.
2. Find the change of the radius.

Express your results through Young's modulus E and Poisson ratio σ of the cylinder.

Problem 2.

A cylinder of radius R and length L is placed inside a very hard tube of the inner radius R and then squeezed by applying a uniform pressure P_0 to its end.

1. Find the change of length.
2. Find the pressure P the cylinder exerts on the tube.
3. What is the maximum possible P at given P_0 ?
4. What is the minimal possible P at given P_0 ? Hint: it's negative.

Express your results through Young's modulus E and Poisson ratio σ of the cylinder.

Homework 15. Due Tuesday, May 5

Problem 1.

A person of mass M is staying at the end of a rigid, thin, and weightless diving board of length L , thickness a , width b , and Young's modulus E . The near end of the board is clamped horizontally. Find the shape of the board.

Problem 2.

A uniform disc of mass M and radius R is hanging horizontally by its center by a vertical thread of radius a , length L and sheer modulus μ . It was given a small initial spin around the vertical direction. What is the period of oscillations? The disc remains horizontal and the thread vertical during the motion.